



**Protocol DNA Purification  
Bacteria**

**with <sup>4</sup>ward-NA® puriSorb®**

**- <sup>4</sup>ward-NA® puriPlate96 -**

**Cat.No. 4w03.P01  
Cat.No. 4w03.P04  
Cat. No. 4w03.P10**

**Version 1.3**

**For research use only**

### Principle

**4ward-NA® DNA purification kits (Bacteria)** are designed for extraction and purification of DNA from a wide range of bacteria species. **4ward-NA® puriSorb® technology** allows the purification in one single step after lysis. Impurities are bound and the purified DNA is recovered in the eluate. The excellent purity of the DNA allows immediate use in all common downstream applications (e.g. PCR).

### Kit contents

The kit contains all necessary reagents for lysis and DNA purification.

Component	Art. No. 4w03.P01	Art. No. 4w03.P04	Art. No. 4w03.P10*
Activation Solution (ActivS)	40 mL	155 mL	400 mL
Buffer 4wC (B)	10 mL	40 mL	100 mL
SDS Solution	10 mL	40 mL	100 mL
RNase A	2.2 mL	9 mL	22 mL
Proteinase K	1.15 mL	4.5 mL	11.5 mL
EDTA	0.25 mL	0.9 mL	2.5 mL
DTT (1,4-Dithio-DL-threitol)	0.3 mL	1.2 mL	2.8 mL
Lysozyme (25 mg/mL)**	30 mg**	112.5 mg**	280 mg**
<b>4ward-NA® puriPlate96</b>	1	4	10
Deep well plate (1.2 mL)	3	12	30
Sealing tapes (plastic)	3	12	30
Storage sealing tapes (alu)	2	8	20

\*other kit sizes on request

\*\*add H<sub>2</sub>O

### Storage Conditions

During shipment all kit components are stable at room temperature. After arrival, store chemicals at **+2 °C to +8 °C**. All other components can be stored at room temperature. Please see expiration date of the kit.

### Safety Information

Proteinase K	Danger	H334 P304+P341, P342+P311
DTT	Warning	H315, H319 P280, P305+P351+P338, P321, P362, P332+P313, P337+P313
SDS	Danger	H315, H318 P305+P351+P338

### Hazard Statements

H315	Causes skin irritation.
H318	Causes serious eye damage.
H319	Causes serious eye irritation.
H334	May cause allergy or asthma symptoms or breathing difficulties if inhaled.

### Precautionary Statements

P280	Wear protective gloves/ protective clothing/ eye protection/ face protection.
P342+P311	If experiencing respiratory symptoms: Call a POISON CENTER or doctor/ physician.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P304+P341	IN INHALED: if breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing.
P321	Specific treatment (see on this label).
P362	Take off contaminated clothing and wash before reuse.
P332+P313	If skin irritation occurs: Get medical advice/ attention.
P337+P313	If eye irritation persists: Get medical advice/ attention.

When working with chemicals, always wear a suitable lab coat, disposable gloves and protective goggles. For more information, please contact us for the appropriate material safety data sheets (MSDS).

### Before starting

#### Activation of puriSorb® in 4ward-NA® puriPlate96

A1	Add 350 µL Activation Solution onto each well of a 4ward-NA® puriPlate96. Incubate for at least 5 min at room temperature. Centrifuge at <u>max. 350 × g</u> for 1 min or apply vacuum for 1 min to remove excess Activation Solution. Discard the deep well plate.
A2	The activated 4ward-NA® puriPlate96 is ready for purification.

- **Preheat an incubator to 56 °C.**
- **Dissolve Lysozyme**  
Add 1.2 mL (4w03.P01), 4.5 mL (4w03.P04) or 11.2 mL (4w03.P10) of sterile water to the lyophilized lysozyme (final concentration is 25 mg/mL).

### Protocol

#### Lysis of bacteria for DNA purification

1. Transfer 0.5 mL of bacterial culture (up to 1.5 OD600) into each well of a deep well plate (1.2 mL). Centrifuge cell suspensions (2,000 × g, 10 min), discard the supernatant.
2. Add 90 µL Buffer 4wC (B), 10 µL Lysozyme and 20 µL RNase A to the cell pellet. Resuspend cells and close the deep well plate (1.2 mL) using a Storage sealing tape. Incubate (56 °C, 20 min).
3. Add 90 µL SDS Solution, 10 µL Proteinase K, 2.5 µL DTT and 2 µL EDTA to each sample. Incubate (56 °C, 60 min).

#### Purification of DNA using 4ward-NA® puriSorb® (ONE-STEP)

P1	Transfer between 60 µL and 120 µL of the lysate into an activated well of the 4ward-NA® puriPlate96. Incubate for at least 3 min and centrifuge at <u>max. 700 × g</u> for 1 min.
	The eluate contains the ready-to-use purified DNA.

### Troubleshooting, FAQ and Special Applications

Please contact us for further information about our innovative DNA Purification System using the **4ward-NA® puriSorb® technology**.

### Contact Information

Web: [www.4ward-na.com](http://www.4ward-na.com)  
Email: [service@4ward-na.com](mailto:service@4ward-na.com)  
Tel: +49 175 948 33 46

### Ordering Information and Prices

For ordering information and prices please contact: [sales@4ward-na.com](mailto:sales@4ward-na.com).

