



**Protocol DNA Purification**

**Plant**

**with 4ward-NA® puriSorb®**

**-4ward-NA® puriColumns -**

**Cat.No. 4w01.010**

**Cat.No. 4w01.050**

**Cat.No. 4w01.250**

**Version 1.3**

**For research use only**

## Principle

**4ward-NA® DNA purification kits (Plant)** are designed for extraction and purification of DNA from a wide range of plant species. **4ward-NA® puriSorb® technology** allows the purification in one single step after lysis. Impurities are bound and the purified DNA is recovered in the eluate. The excellent purity of the DNA allows immediate use in all common downstream applications (e.g. PCR).

## Kit contents

The kit contains all necessary reagents for lysis and DNA purification.

Component	Art. No. 4w01.010	Art. No. 4w01.050	Art. No. 4w01.250
Activation Solution (ActivS)	4.5 mL	20.5 mL	100 mL
Buffer 4wA (F)	3.5 mL	16.5 mL	81 mL
Buffer 4wB (H)	2.5 mL	11 mL	54 mL
Protease	0.25 mL	1.2 mL	6 mL
SDS	0.15 mL	0.6 mL	2.9 mL
RNase A	on request	on request	on request
DTT (1,4-Dithio-DL-threitol)	on request	on request	on request
4ward-NA® puriColumns	10	50	250
Waste Tubes	10	50	250
DNA Tubes	10	50	250

## Storage Conditions

During shipment all kit components are stable at room temperature. After arrival, store chemicals at **+2 °C to +8 °C**. All other components can be stored at room temperature. Please see expiration date of the kit.

**Safety Information**

Protease	Danger	H315, H318, H334, H335 P261, P280, P342+P311, P305+P351+P338
DTT	Warning	H315, H319 P280, P305+P351+P338, P321, P362, P332+P313, P337+P313
SDS	Danger	H315, H318 P305+P351+P338

**Hazard Statements**

H315	Causes skin irritation.
H318	Causes serious eye damage.
H319	Causes serious eye irritation.
H334	May cause allergy or asthma symptoms or breathing difficulties if inhaled.
H335	May cause respiratory irritation.

**Precautionary Statements**

P261	Avoid breathing dust/ fume/ gas/ mist/ vapors/ spray.
P280	Wear protective gloves/ protective clothing/ eye protection/ face protection.
P342+P311	If experiencing respiratory symptoms: Call a POISON CENTER or doctor/ physician.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P321	Specific treatment (see on this label).
P362	Take off contaminated clothing and wash before reuse.
P332+P313	If skin irritation occurs: Get medical advice/ attention.
P337+P313	If eye irritation persists: Get medical advice/ attention.

When working with chemicals, always wear a suitable lab coat, disposable gloves and protective goggles. For more information, please contact us for the appropriate material safety data sheets (MSDS).

## Before starting

### Activation of puriSorb® in 4ward-NA® puriColumns

A1	Add <b>350 µL Activation Solution</b> onto each well of a <b>4ward-NA® puriColumn</b> . Incubate for at least 5 min at room temperature. Centrifuge at <u>max. 350 × g</u> for <b>1 min</b> to remove excess Activation Solution. Discard the Waste Tube.
A2	The activated <b>4ward-NA® puriColumn</b> is ready for purification.

- Preheat a thermomixer to 60 °C.

## Protocol

### Lysis of plant tissue for DNA purification

1. Transfer **5 pieces of leaf tissue** (e.g. 5 × 5 mm) into a 1.5 mL or 2.0 mL reaction tube.
2. Add **280 µL Buffer 4wA (F)** and **20 µL Protease**.
3. Incubate (**60 °C, 30 min up to overnight**).

### Lysis of homogenized plant tissue for DNA purification

1. Homogenize/grind up to **50 mg of plant tissue** with **185 µL Buffer 4wB (H)**  
(optional: add **3 µL DTT** / **12 µL RNase A**).
2. Add **70 µL Buffer 4wA (F)**, **20 µL Protease** and **10 µL SDS**.
3. Incubate (**60 °C, 30 min up to overnight**), if necessary, centrifuge for a clear supernatant for DNA purification.

### Lysis of plant seeds for DNA purification

1. Homogenize/grind up to **30 mg of plant seeds** or use up to 30 mg plant seed meal.
2. Add **280 µL Buffer 4wA (F)** and **20 µL Protease** (optional: add **3 µL DTT**).
3. Incubate (**60 °C, 30 min up to overnight**).

### Purification of DNA using 4ward-NA® puriSorb® (ONE-STEP)

P1	Transfer between <b>60 µL</b> and <b>120 µL</b> of the lysate to an activated <b>4ward-NA® puriColumn</b> . Incubate for at least <b>3 min</b> and centrifuge at <u>max. 700 × g</u> for <b>1 min</b> .
	The eluate contains the ready-to-use purified DNA.

**Troubleshooting, FAQ and Special Applications**

Please contact us for further information about our innovative DNA Purification System using the **4ward-NA® puriSorb®** technology.

**Contact Information**

Web: [www.4ward-na.com](http://www.4ward-na.com)  
Email: [service@4ward-na.com](mailto:service@4ward-na.com)  
Tel: +49 175 948 33 46

**Ordering Information and Prices**

For ordering information and prices please contact: [sales@4ward-na.com](mailto:sales@4ward-na.com).

